Sign in

Google

Web Images Video News Maps more » Search Advanced Search mcf cells irs Preferences

Web

Results 1 - 10 of about 111,000 for mcf cells irs. (0.30 seconds)

Retinoic acid-induced growth arrest of MCF-7 cells involves the ...

Treatment of MCF-7 cells with RA caused a significant reduction in IRS-1 ... RA resistance was observed in MCF-7 cells overexpressing IRS-1 but not IGF-IR. ... www.nature.com/onc/journal/v22/n22/abs/1206485a.html - Similar pages

Progesterone crosstalks with insulin-like growth factor signaling ... Immunoblot analysis of total cell lysates demonstrated that a single dose of 10-8 M R5020 for 24 h remarkably increased IRS-2 levels in MCF-7 cells, ... www.nature.com/onc/journal/v22/n44/full/1206803a.html - Similar pages

Insulin-like growth factor 1 and oestradiol promote cell ...

However, IGF-1 induced tyrosine phosphorylation of IGF-1R was enhanced by E2 treatment of the MCF-7 cells. IRS-1 protein values were increased by E2, ... mp.bmjjournals.com/cgi/content/full/54/3/149 - Similar pages

[PDF] Insulin-like growth factor 1 and oestradiol promote cell ...

File Format: PDF/Adobe Acrobat synchronise the cells in G0 phase. However,. IGF-1 induced tyrosine phosphorylation of. IGF-1R was enhanced by E2 treatment of the. MCF-7 cells. IRS-1 ... mp.bmjjournals.com/cgi/reprint/54/3/149.pdf - Similar pages

Retinoic acid-induced growth arrest of MCF-7 cells involves the ...

RA resistance was observed in MCF-7 cells overexpressing IRS-1 but not IGF-IR. This suggests that RA-mediated growth inhibition requires the selective ... www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve& db=PubMed&list_uids=12776186&dopt=Abstract - Similar pages

Motility response to insulin-like growth factor-I (IGF-I) in MCF-7 ...

In MCF-7/ATCC cells, IGF-I stimulation was associated with a movement of IRS-2 to the leading edge of filopodia. Thus, patterns of integrin expression among ... www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve& db=PubMed&list_uids=14997047&dopt=Citation - Similar pages [More results from www.ncbi.nlm.nih.gov]

<u>Differential roles of IRS-1 and SHC signaling pathways in breast ...</u>

In MCF-7 cells, down-regulation of IRs-I by 80-85% strongly suppressed anchoragedependent and-independent growth and induced apoptotic cell death under ... cat.inist.fr/?aModele=afficheN&cpsidt=2792077 - Similar pages

CAT.INIST.FR

Motility response to insulin-like growth factor-I (IGF-I) in MCF-7 cells is associated with IRS-2 activation and integrin expression ... cat.inist.fr/?aModele=afficheN&cpsidt=15433162 - Similar pages [More results from cat.inist.fr]

IRS Cancer Connection -- Gough 2006 (365): tw417 -- Science's STKE Dearth et al. used transgenic mice forced to overexpress IRS-1 or IRS-2, or an immortalized but nontumorigenic cell line (MCF-10A) also forced to ... stke.sciencemag.org/cgi/content/abstract/sigtrans;2006/365/tw417 - Similar pages

Overexpression of insulin receptor substrate 1 (IRS-1) in the ... We report here that in **MCF-7 cells** overexpressing **IRS-1**, estrogen requirements for growth in monolayer culture as well as in soft agar were reduced. ... clincancerres.aacrjournals.org/cgi/content/abstract/1/11/1429 - <u>Similar pages</u>

Result Page:	1 2 3 4 5 6 7 8 9 10	Next
mcf cells Search within results Lan	s irs Suage Tools Search Tips Dis	Search satisfied? Help us improve

Google Home - Advertising Programs - Business Solutions - About Google ©2007 Google